antibodies -online.com





anti-STAG2 antibody (AA 41-150) (Biotin)



Go to Product page

()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL	
Target:	STAG2	
Binding Specificity:	AA 41-150	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This STAG2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SA2	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	STAG2	
Alternative Name:	SA2 (STAG2 Products)	
Background:	Synonyms: Cohesin Subunit SA 2, Cohesin subunit SA-2, DKFZp686P168, DKFZp781H1753, SA	

2, SA2, SCC3 homolog 2, STAG 2, STAG 2, stag2, STAG2_HUMAN, Stromal antigen 2,	
FLJ25871. bA51701.1.	

Background: Component of cohesin complex, a complex required for the cohesion of sister chromatids after DNA replication. The cohesin complex apparently forms a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. The cohesin complex may also play a role in spindle pole assembly during mitosis.

Gene ID: 10735

Pathways: Stem Cell Maintenance

Application Details

Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months