# antibodies -online.com





# anti-SKA1 antibody (AA 181-255) (Alexa Fluor 647)



( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Quantity:	100 μL	
Target:	SKA1	
Binding Specificity:	AA 181-255	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SKA1 antibody is conjugated to Alexa Fluor 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SKA1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	SKA1
Alternative Name:	SKA1 (SKA1 Products)
Background: Synonyms: Ska1, SKA1_HUMAN, Spindle and kinetochore associated complex subunit	

Spindle and kinetochore associated protein 1, Spindle and kinetochore-associated protein 1, Spindle and KT kinetochore associated 1, Spindle and KT associated 1, C18orf24, MGC10200. Background: Component of the SKA1 complex, a microtubule-binding subcomplex of the outer kinetochore that is essential for proper chromosome segregation. Required for timely anaphase onset during mitosis, when chromosomes undergo bipolar attachment on spindle microtubules leading to silencing of the spindle checkpoint. The SKA1 complex is a direct component of the kinetochore-microtubule interface and directly associates with microtubules as oligomeric assemblies. The complex facilitates the processive movement of microspheres along a microtubule in a depolymerization-coupled manner. In the complex, it mediates the interaction with microtubules.

Pathways:

M Phase

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months