

Datasheet for ABIN915462

anti-SLC16A7 antibody (AbBy Fluor® 647)



Go to Product page

_				
	۱۱ / ۱	rv		۱۸/
	' V '	 ı v	Ι.	v v

Quantity:	100 μL	
Target:	SLC16A7	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SLC16A7 antibody is conjugated to AbBy Fluor® 647	
Application:	Western Blotting (WB)	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLC16A7	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	SLC16A7
Alternative Name:	Slc16a7/Mct2 (SLC16A7 Products)
Background:	Synonyms: MCT 2, MCT, Monocarboxylate transporter 2, MonocarboxylateTransporter 2,
	MOT2, MOT2_HUMAN, SLC16A7, Solute carrier family 16 member 7, Solute carrier Family 16

Monocarboxylic Acid Transporters Member 7.

Background: Tissues with few or no mitochondria, such as erythrocytes and tumor cells, depend largely on glycolysis to generate ATP. The major end products of glycolysis, pyruvate and lactate, must be eliminated from these cells to enable continued glycolytic flux and prevent toxic effects. H+/monocarboxylate transporters (MCTs) mediate the transport of lactate and pyruvate. Human MCT2 has a high affinity for the transport of pyruvate (summary by Lin et al., 1998 [PubMed 9786900]).[supplied by OMIM, Feb 2011].

Gene ID:

Expiry Date:

9194

12 months

Application Details

Application Notes:	IF(IHC-P) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	