antibodies -online.com





anti-SMURF2 antibody (AA 601-700) (Alexa Fluor 555)



_			
	$ \backslash / \cap$	r\/I	ew
\sim	' V C	. I V I	C V V

Quantity:	100 μL
Target:	SMURF2
Binding Specificity:	AA 601-700
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SMURF2 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SMURF2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	SMURF2
Alternative Name:	SMURF2 (SMURF2 Products)

Target Details

Background:	Synonyms: E3 ubiquitin-protein ligase SMURF2, hSMURF2, SMAD ubiquitination regulatory		
	factor 2, SMAD-specific E3 ubiquitin-protein ligase 2, SMURF2		
	Background: E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-		
	conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to		
	targeted substrates. Interacts with SMAD1 and SMAD7 in order to trigger their ubiquitination		
	and proteasome-dependent degradation. In addition, interaction with SMAD7 activates		
	autocatalytic degradation, which is prevented by interaction with SCYE1. Forms a stable		
	complex with the TGF-beta receptor-mediated phosphorylated SMAD2 and SMAD3. In this way,		
	SMAD2 may recruit substrates, such as SNON, for ubiquitin-mediated degradation. Enhances		
	the inhibitory activity of SMAD7 and reduces the transcriptional activity of SMAD2.		
	Coexpression of SMURF2 with SMAD1 results in considerable decrease in steady-state level of		
	SMAD1 protein and a smaller decrease of SMAD2 level.		
Gene ID:	64750		
UniProt:	Q9HAU4		
Application Details			
Application Notes:	FCM 1:20-100		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

		1.	
\vdash	land	ling	
1 1	ıaııu	11111	۰

Expiry Date:

12 months