antibodies -online.com





anti-Sortilin 1 antibody (AA 731-831) (Cy7)



Overview

Quantity:	100 μL
Target:	Sortilin 1 (SORT1)
Binding Specificity:	AA 731-831
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Sortilin 1 antibody is conjugated to Cy7
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Sortilin
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	Sortilin 1 (SORT1)
Alternative Name:	Sortilin (SORT1 Products)

Target Details

Background:	Synonyms: SORT1,100 kDa NT receptor, Glycoprotein 95 Short name: Gp95, Neurotensin
	receptor 3 Short name: NT3 Short name: NTR3, SORT_HUMAN
	Background: Functions as a sorting receptor in the Golgi compartment and as a clearance
	receptor on the cell surface. Required for protein transport from the Golgi apparatus to the
	lysosomes by a pathway that is independent of the mannose-6-phosphate receptor (M6PR).
	Also required for protein transport from the Golgi apparatus to the endosomes. Promotes
	neuronal apoptosis by mediating endocytosis of the proapoptotic precursor forms of BDNF
	(proBDNF) and NGFB (proNGFB). Also acts as a receptor for neurotensin. May promote
	mineralization of the extracellular matrix during osteogenic differentiation by scavenging
	extracellular LPL. Probably required in adipocytes for the formation of specialized storage
	vesicles containing the glucose transporter SLC2A4/GLUT4 (GLUT4 storage vesicles, or GSVs).
	These vesicles provide a stable pool of SLC2A4 and confer increased responsiveness to insulin
	May also mediate transport from the endoplasmic reticulum to the Golgi.
Gene ID:	6272
UniProt:	Q99523
Pathways:	Neurotrophin Signaling Pathway
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Buffer: Preservative:	
	50 % Glycerol.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months