# antibodies .- online.com







## anti-Sortilin 1 antibody (AA 731-831) (Alexa Fluor 350)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	Sortilin 1 (SORT1)	
Binding Specificity:	AA 731-831	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Sortilin 1 antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Sortilin	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Guinea Pig	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	Sortilin 1 (SORT1)
Alternative Name:	Sortilin (SORT1 Products)

#### Target Details

Background:	Synonyms: SORT1,100 kDa NT receptor, Glycoprotein 95 Short name: Gp95, Neurotensin	
	receptor 3 Short name: NT3 Short name: NTR3, SORT_HUMAN	
	Background: Functions as a sorting receptor in the Golgi compartment and as a clearance	
	receptor on the cell surface. Required for protein transport from the Golgi apparatus to the	
	lysosomes by a pathway that is independent of the mannose-6-phosphate receptor (M6PR).	
	Also required for protein transport from the Golgi apparatus to the endosomes. Promotes	
	neuronal apoptosis by mediating endocytosis of the proapoptotic precursor forms of BDNF	
	(proBDNF) and NGFB (proNGFB). Also acts as a receptor for neurotensin. May promote	
	mineralization of the extracellular matrix during osteogenic differentiation by scavenging	
	extracellular LPL. Probably required in adipocytes for the formation of specialized storage	
	vesicles containing the glucose transporter SLC2A4/GLUT4 (GLUT4 storage vesicles, or GSVs).	
	These vesicles provide a stable pool of SLC2A4 and confer increased responsiveness to insulin	
	May also mediate transport from the endoplasmic reticulum to the Golgi.	
Gene ID:	6272	
UniProt:	Q99523	
Pathways:	Neurotrophin Signaling Pathway	
Application Details		
Application Notes:	FCM 1:20-100	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Buffer: Preservative:		
	50 % Glycerol.	

### Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months