antibodies -online.com





anti-Sortilin 1 antibody (AA 731-831) (Alexa Fluor 488)



Go to Product page

\sim				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL	
Target:	Sortilin 1 (SORT1)	
Binding Specificity:	AA 731-831	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Sortilin 1 antibody is conjugated to Alexa Fluor 488	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Sortilin	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Guinea Pig	
Purification:	Purified by Protein A.	

Target Details

Target:	Sortilin 1 (SORT1)	
Alternative Name:	Sortilin (SORT1 Products)	

Target Details

Background:	Synonyms: SORT1,100 kDa NT receptor, Glycoprotein 95 Short name: Gp95, Neurotensin		
	receptor 3 Short name: NT3 Short name: NTR3, SORT_HUMAN		
	Background: Functions as a sorting receptor in the Golgi compartment and as a clearance		
	receptor on the cell surface. Required for protein transport from the Golgi apparatus to the		
	lysosomes by a pathway that is independent of the mannose-6-phosphate receptor (M6PR).		
	Also required for protein transport from the Golgi apparatus to the endosomes. Promotes		
	neuronal apoptosis by mediating endocytosis of the proapoptotic precursor forms of BDNF		
	(proBDNF) and NGFB (proNGFB). Also acts as a receptor for neurotensin. May promote		
	mineralization of the extracellular matrix during osteogenic differentiation by scavenging		
	extracellular LPL. Probably required in adipocytes for the formation of specialized storage		
	vesicles containing the glucose transporter SLC2A4/GLUT4 (GLUT4 storage vesicles, or GSVs).		
	These vesicles provide a stable pool of SLC2A4 and confer increased responsiveness to insulin		
	May also mediate transport from the endoplasmic reticulum to the Golgi.		
Gene ID:	6272		
UniProt:	Q99523		
Pathways:	Neurotrophin Signaling Pathway		
Application Details			
Application Notes:	FCM 1:20-100		
	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.		
Buffer: Preservative:			

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months