antibodies -online.com





anti-STK40 antibody (AA 231-330) (HRP)



()	1 /	\sim	KI /	110	Νę
	1//	\vdash	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	STK40	
Binding Specificity:	AA 231-330	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This STK40 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human STK40	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	STK40
Alternative Name:	STK40 (STK40 Products)

Target Details

rarget betails		
Background:	Synonyms: Ser/Thr like kinase, Serine/threonine kinase 40, Serine/threonine protein kinase 40, SgK495, SHIK, SINK homologous serine/threonine protein kinase, SINK-homologous serine/threonine-protein kinase, STK 40, Sugen kinase 495, STK40_HUMAN.	
	Background: STK40 may be a negative regulator of NF-kappa-B and p53-mediated gene transcription. There are four named isoforms.	
Gene ID:	83931	
UniProt:	Q8N2I9	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	