

Datasheet for ABIN917377

anti-TFAP2C antibody (AA 2-100) (AbBy Fluor® 555)



Go to Product page

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Target:

Alternative Name:

Quantity:	100 μL
Target:	TFAP2C
Binding Specificity:	AA 2-100
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TFAP2C antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence
	(Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human TFAP2C
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Rabbit
Purification:	Purified by Protein A.
Target Details	

TFAP2C

AP2 gamma (TFAP2C Products)

Target Details

Synonyms: ERF1, TFAP2G, hAP-2g, AP2-GAMMA, Transcription factor AP-2 gamma, Activating	
enhancer-binding protein 2 gamma, Transcription factor ERF-1, TFAP2C	
Background: Sequence-specific DNA-binding protein that interacts with inducible viral and	
cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the	
consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of	
important biological functions including proper eye, face, body wall, limb and neural tube	
development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha	
and MYC. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast	
cancer.	
7022	
Q92754	
Stem Cell Maintenance	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
handled by trained staff only.	
-20 °C	
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
12 months	