

## Datasheet for ABIN917380

## anti-TFAP2C antibody (AA 2-100) (AbBy Fluor® 350)



Go to Product page

_					
	1//	r	Vİ	$\triangle$	۸/
	V		VI		/ V

Quantity:	100 μL	
Target:	TFAP2C	
Binding Specificity:	AA 2-100	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TFAP2C antibody is conjugated to AbBy Fluor® 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human TFAP2C	

Immunogen:	KLH conjugated synthetic peptide derived from human TFAP2C
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	TFAP2C
Alternative Name:	AP2 gamma (TFAP2C Products)

## **Target Details**

Background:	Synonyms: ERF1, TFAP2G, hAP-2g, AP2-GAMMA, Transcription factor AP-2 gamma, Activating		
	enhancer-binding protein 2 gamma, Transcription factor ERF-1, TFAP2C		
	Background: Sequence-specific DNA-binding protein that interacts with inducible viral and		
	cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the		
	consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of		
	important biological functions including proper eye, face, body wall, limb and neural tube		
	development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha		
	and MYC. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast		
	cancer.		
Gene ID:	7022		
UniProt:	Q92754		
Pathways:	Stem Cell Maintenance		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		