# antibodies .- online.com







# anti-TNIP2 antibody (AA 85-180) (Alexa Fluor 555)



| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | TNIP2  |  |
| Binding Specificity: | AA 85-180  |  |
| Reactivity:          | Human, Rat   |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This TNIP2 antibody is conjugated to Alexa Fluor 555   |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human TNIP2/ABIN2 |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Cross-Reactivity:     | Human, Rat  |  |
| Predicted Reactivity: | Mouse,Dog,Horse   |  |
| Purification:         | Purified by Protein A.  |  |

## Target Details

| Target:           | TNIP2                  |
|-------------------|------------------------|
| Alternative Name: | TNIP2 (TNIP2 Products) |

## **Target Details**

| •                   |   |  |
|---------------------|---|--|
| Background:         | Synonyms: KLIP, ABIN2, FLIP1, TNFAIP3-interacting protein 2, A20-binding inhibitor of NF-   |  |
|                     | kappa-B activation 2, ABIN-2, Fetal liver LKB1-interacting protein, TNIP2                   |  |
|                     | Background: Inhibits NF-kappa-B activation by blocking the interaction of RIPK1 with its    |  |
|                     | downstream effector NEMO/IKBKG. Forms a ternary complex with NFKB1 and MAP3K8 but           |  |
|                     | appears to function upstream of MAP3K8 in the TLR4 signaling pathway that regulates         |  |
|                     | MAP3K8 activation. Involved in activation of the MEK/ERK signaling pathway during innate    |  |
|                     | immune response, this function seems to be stimulus- and cell type specific. Required for   |  |
|                     | stability of MAP3K8. Involved in regulation of apoptosis in endothelial cells, promotes TEK |  |
|                     | agonist-stimulated endothelial survival. May act as transcriptional coactivator when        |  |
|                     | translocated to the nucleus. Enhances CHUK-mediated NF-kappa-B activation involving NF-     |  |
|                     | kappa-B p50-p65 and p50-c-Rel complexes.  |  |
| Gene ID:            | 79155   |  |
| UniProt:            | Q8NFZ5  |  |
| Pathways:           | Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin     |  |
| Application Details |   |  |
| Application Notes:  | IF(IHC-P) 1:50-200  |  |
|                     | IF(IHC-F) 1:50-200  |  |
|                     | IF(ICC) 1:50-200  |  |
| Restrictions:       | For Research Use only   |  |
| Handling            |   |  |
| Format:             | Liquid  |  |
| Concentration:      | 1 μg/μL   |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |  |
|                     | 50 % Glycerol.  |  |
| Preservative:       | ProClin   |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be         |  |
|                     | handled by trained staff only.  |  |
| Storage:            | -20 °C  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.           |  |
|                     |   |  |

| $\vdash$ | land  | linc  |
|----------|-------|-------|
|          | iaria | 11116 |

Expiry Date:

12 months