.-online.com antibodies

Datasheet for ABIN918413 anti-NR2C2AP antibody (AA 81-139) (Alexa Fluor 488)



Overview

Quantity:	100 µL
Target:	NR2C2AP
Binding Specificity:	AA 81-139
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR2C2AP antibody is conjugated to Alexa Fluor 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human of human TRA16
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	NR2C2AP
Alternative Name:	TRA16 (NR2C2AP Products)
Background:	Synonyms: NR2C2AP, Nuclear receptor 2C2 associated protein, Repressor for TR4

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN918413 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details

associated protein TRA16, Testicular orphan nuclear receptor-4 TR4-associated pr NR2CA_HUMAN. Background: TRA16 may act as a repressor of NR2C2-mediated transactivation by the binding between NR2C2/TR4 and the TR4-response element in target genes. Gene ID: 126382 Pathways: Nuclear Receptor Transcription Pathway Application Details IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling IF(ICC) 1:50-200 Preservative: ProClin Preservative: ProClin Preservative: ProClin Preca		
NR2CA_HUMAN. Background: TRA16 may act as a repressor of NR2C2-mediated transactivation by the binding between NR2C2/TR4 and the TR4-response element in target genes.Gene ID:126382Pathways:Nuclear Receptor Transcription PathwayApplication DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingIuquidFormat:LiquidConcentration:1 µg/µLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Giycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, while handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		transactivation, TR4 orphan receptor associated 16 kDa protein, TR4 orphan receptor
Background: TRA16 may act as a repressor of NR2C2-mediated transactivation by the binding between NR2C2/TR4 and the TR4-response element in target genes.Gene ID:126382Pathways:Nuclear Receptor Transcription PathwayApplication DetailsIF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200Restrictions:For Research Use onlyHandlingIuquidConcentration:1 µg/µLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a PDISONOUS AND HAZARDOUS SUBSTANCE, whit handled by trained staff only.Storage Comment::20 °C		associated protein TRA16, Testicular orphan nuclear receptor-4 TR4-associated protein, NR2CA_HUMAN.
the binding between NR2C2/TR4 and the TR4-response element in target genes. Gene ID: 126382 Pathways: Nuclear Receptor Transcription Pathway Application Details IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Iquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whit handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Gene ID: 126382 Pathways: Nuclear Receptor Transcription Pathway Application Details IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whit handled by trained staff only. Storage -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		Background: TRA16 may act as a repressor of NR2C2-mediated transactivation by suppressing
Pathways: Nuclear Receptor Transcription Pathway Application Details Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL. Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whith handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		the binding between NR2C2/TR4 and the TR4-response element in target genes.
Application Details Application Notes: IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, while handled by trained staff only. Storage: -20 °C Storage Comment: Storage Comment:	Gene ID:	126382
Application Notes: IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Iuquid Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whit handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Pathways:	Nuclear Receptor Transcription Pathway
IF(IHC-F) 1:50-200Restrictions:For Research Use onlyHandlingFormat:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, while handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Details	
IF(ICC) 1:50-200 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, while handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Notes:	IF(IHC-P) 1:50-200
Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whith handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		IF(IHC-F) 1:50-200
HandlingFormat:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		IF(ICC) 1:50-200
Format:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Restrictions:	For Research Use only
Concentration:1 µg/µLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Handling	
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Pr 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Format:	Liquid
50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Concentration:	1 µg/µL
Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only.Storage:-20 °CStorage Comment:Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
handled by trained staff only. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Preservative:	ProClin
Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		handled by trained staff only.
	Storage:	-20 °C
Expiny Date: 12 months	Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
	Expiry Date:	12 months