antibodies .- online.com







anti-TSPYL5 antibody (AA 251-350) (Cy7)



| () | 11/ | IN | /ie | A . |
|-----|-------------------|------|--------|--------|
| | / // | ۱ ات | / (− | ' \/\/ |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | TSPYL5 | |
| Binding Specificity: | AA 251-350 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This TSPYL5 antibody is conjugated to Cy7 | |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human TSPYL5 |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | TSPYL5 |
|-------------------|--------------------------|
| Alternative Name: | TSPYL5 (TSPYL5 Products) |

Target Details

Expiry Date:

12 months

| rarget betails | | |
|---------------------|---|--|
| Background: | Synonyms: Testis-specific Y-encoded-like protein 5, TSPY-like protein 5, TSPYL5, KIAA1750 Background: Involved in modulation of cell growth and cellular response to gamma radiation probably via regulation of the Akt signaling pathway. Involved in regulation of p53/TP53. Suppresses p53/TP53 protein levels and promotes its ubiquitination, the function is dependent on USP7 and independent on MDM2. Proposed to displace p53/TP53 from interaction with USP7. | |
| Gene ID: | 85453 | |
| UniProt: | Q86VY4 | |
| Application Details | | |
| Application Notes: | FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| | | |