

Datasheet for ABIN919270

anti-UGDH antibody (AA 421-494) (AbBy Fluor® 488)



Go to Product page

Overview	
Quantity:	100 μL
Target:	UGDH
Binding Specificity:	AA 421-494
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UGDH antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human UGDH/UDPGDH
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	UGDH
Alternative Name:	UGDH (UGDH Products)

Target Details

Background:	Synonyms: UDP glucose dehydrogenase, GDH, UDP Glc dehydrogenasev UDP GlcDH, UDP		
	glucose 6 dehydrogenase, UDP-Glc dehydrogenase, UDP-GlcDH, UDP-glucose 6-		
	dehydrogenase, UDPGDH, UGD, Ugdh, UGDH_HUMAN, Uridine diphospho glucose		
	dehydrogenase.		
	Background: The protein encoded by this gene converts UDP-glucose to UDP-glucuronate and		
	thereby participates in the biosynthesis of glycosaminoglycans such as hyaluronan, chondroitin sulfate, and heparan sulfate. These glycosylated compounds are common components of the extracellular matrix and likely play roles in signal transduction, cell migration, and cancer growth and metastasis. The expression of this gene is up-regulated by transforming growth factor beta and down-regulated by hypoxia. Alternative splicing results in multiple transcript		
		variants.[provided by RefSeq, May 2010].	
		Gene ID:	7358
		Pathways:	Glycosaminoglycan Metabolic Process
		Application Details	
	Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		