

## Datasheet for ABIN919272

## anti-UGDH antibody (AA 421-494) (AbBy Fluor® 555)



_				
( )	ve.	rv/	101	Λ

Overview		
Quantity:	100 μL	
Target:	UGDH	
Binding Specificity:	AA 421-494	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UGDH antibody is conjugated to AbBy Fluor® 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human UGDH/UDPGDH	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	UGDH	
Alternative Name:	UGDH (UGDH Products)	

## **Target Details**

Background:	Synonyms: UDP glucose dehydrogenase, GDH, UDP Glc dehydrogenasev UDP GlcDH, UDP	
	glucose 6 dehydrogenase, UDP-Glc dehydrogenase, UDP-GlcDH, UDP-glucose 6-	
	dehydrogenase, UDPGDH, UGD, Ugdh, UGDH_HUMAN, Uridine diphospho glucose	
	dehydrogenase.	
	Background: The protein encoded by this gene converts UDP-glucose to UDP-glucuronate and	
	thereby participates in the biosynthesis of glycosaminoglycans such as hyaluronan, chondroitin	
	sulfate, and heparan sulfate. These glycosylated compounds are common components of the	
	extracellular matrix and likely play roles in signal transduction, cell migration, and cancer growth	
	and metastasis. The expression of this gene is up-regulated by transforming growth factor beta	
	and down-regulated by hypoxia. Alternative splicing results in multiple transcript	
	variants.[provided by RefSeq, May 2010].	
Gene ID:	7358	
Pathways:	Glycosaminoglycan Metabolic Process	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	