# antibodies .- online.com







# anti-STRAP antibody (AA 51-150) (Alexa Fluor 647)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	STRAP
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STRAP antibody is conjugated to Alexa Fluor 647
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Unrip
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	STRAP
Alternative Name:	Unrip (STRAP Products)
Background:	Synonyms: MAP activator with WD repeats, MAWD, PTWD, serine/threonine kinase receptor

associated protein, STRAP, UNR-interacting protein, UNRIP, WD 40 repeat protein PT WD, STRAP\_HUMAN.

Background: Unrip is part of the SMN complex that plays a role in snRNP assembly in the cytoplasm and pre mRNA splicing in the nucleus. Unrip interacts directly with Gemin 6 and Gemin 7 in the SMN complex. It is associated with the complex in the cytoplasm but not in the nucleus and thus is thought to play a role in it's subcellular localisation. It is a ubiquitously expressed protein that acts as an inhibitor of TGF-beta signaling and an important regulator of cell proliferation. Stable expression of STRAP results in activation of the mitogen-activated protein kinase/extracellular signal-regulated kinase pathway and in down-regulation of the cyclin-dependent kinase inhibitor p21 (Cip1), which results in retinoblastoma protein hyperphosphorylation. Upregulation of STRAP might play a role in tumor development as an early event for colorectal cancers. It is the first component of the U snRNP assembly machinery that associates with SMN complex in a compartment-specific way and plays a crucial role in the intracellular distribution of SMN1.

Gene ID:

11171

#### **Application Details**

Application Notes:	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

1	1 11	١٠
$\vdash$	land	∥n∩
	ıarıa	

Expiry Date:

12 months