# antibodies -online.com





# anti-WNT6 antibody (AA 281-365) (Alexa Fluor 488)



Go to Product page

| ( ) | 11/0   | K\ /   | iew                 | 1 |
|-----|--------|--------|---------------------|---|
|     | $\cup$ | 'I V/I | $\square \vee \vee$ | ı |
|     |        |        |                     |   |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | WNT6   |
| Binding Specificity: | AA 281-365   |
| Reactivity:          | Human, Mouse   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This WNT6 antibody is conjugated to Alexa Fluor 488  |
| Application:         | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human Wnt6 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human, Mouse   |
| Predicted Reactivity: | Rat  |
| Purification:         | Purified by Protein A.                                   |

## **Target Details**

| Target:           | WNT6                 |
|-------------------|----------------------|
| Alternative Name: | Wnt6 (WNT6 Products) |

## **Target Details**

| Background:         | Synonyms: Protein Wnt-6, WNT6  |  |
|---------------------|--|--|
|                     | Background: Ligand for members of the frizzled family of seven transmembrane receptors.                    |  |
|                     | Probable developmental protein. May be a signaling molecule which affects the development of               |  |
|                     | discrete regions of tissues. Is likely to signal over only few cell diameters. Together with CAV1          |  |
|                     | may promote chemoresistance of gastric cancer cells to DNA-damaging anthracycline drugs                    |  |
|                     | through the activation of the canonical Wnt receptor signaling pathway.                                    |  |
| Gene ID:            | 7475   |  |
| UniProt:            | Q9Y6F9   |  |
| Pathways:           | WNT Signaling, Tube Formation  |  |
| Application Details |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                        |  |
|                     | handled by trained staff only.   |  |
| Storage:            | -20 °C   |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                          |  |
| Expiry Date:        | 12 months  |  |
|                     |  |  |