

Datasheet for ABIN925777

anti-POLR2B antibody (N-Term)





0 -	4.4	Dan decat	
GO	lO	Product	page


()	11 /	er	~\ /	10	1 4
	1 / /	-1	١/	-	1///
\sim	٧.	\sim 1	v	\sim	· v v

Quantity:	100 μg	
Target:	POLR2B	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse, Rat, Dog, Cow, Xenopus laevis, Chicken	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This POLR2B antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	POLR2 B antibody was raised in rabbit using the N terminal of POLR2 as the immunogen	
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Dog (Canine), Cow (Bovine), Frog, Chicken	
Target Details		
Target:	POLR2B	
Alternative Name:	POLR2B (POLR2B Products)	
Background:	POLR2B is the second largest subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. This subunit, in combination with at least two other polymerase subunits, forms a structure within the polymerase that maintains contact in the active site of the enzyme between the DNA template and the newly synthesized RNA. Synonyms: Polyclonal POLR2B antibody, Anti-POLR2B antibody, polymerase, RNA II, DNA	

Target Details

Target Details			
	directed polypeptide B, 140kDa antibody.		
Pathways:	Regulatory RNA Pathways, DNA Damage Repair		
Application Details			
Application Notes:	WB: 5 µg/mL		
	Optimal conditions should be determined by the investigator.		
Comment:	POLR2B Blocking Peptide, catalog no. 33R-9401, is also available for use as a blocking control		
	in assays to test for specificity of this POLR2B antibody		
Restrictions:	For Research Use only		
Handling			
Format:	Lyophilized		
Concentration:	Lot specific		
Buffer:	Lyophilized powder. Add 100 μL of distilled water. Final antibody concentration is 1 mg/mL in		
	PBS buffer.		
Handling Advice:	Avoid repeated freeze/thaw cycles.		
Storage:	4 °C/-20 °C		
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.		

Images



Western Blotting

Image 1. POLR2B antibody (20R-1076) used at 5 ug/ml to detect target protein.