

Datasheet for ABIN925792  
**anti-TRIM17 / RNF16 antibody (C-Term)**



[Go to Product page](#)

1 Image

## Overview

|                      |   |
|----------------------|---|
| Quantity:            | 100 µg  |
| Target:              | TRIM17 / RNF16 (TRIM17)                       |
| Binding Specificity: | C-Term  |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal                                    |
| Conjugate:           | This TRIM17 / RNF16 antibody is un-conjugated |
| Application:         | Western Blotting (WB)                         |

## Product Details

|            |  |
|------------|--|
| Immunogen: | TRIM17 antibody was raised in rabbit using the C terminal of TRIM17 as the immunogen |
|------------|--|

## Target Details

|                   |  |
|-------------------|--|
| Target:           | TRIM17 / RNF16 (TRIM17)  |
| Alternative Name: | TRIM17 ( <a href="#">TRIM17 Products</a> )   |
| Background:       | TRIM17 is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. The protein is expressed almost exclusively in the testis, but its function is unknown. Multiple alternatively spliced transcript variants have been found for this gene. Synonyms: Polyclonal TRIM17 antibody, Anti-TRIM17 antibody, tripartite motif-containing 17 antibody. |


## Application Details

|                    |  |
|--------------------|--|
| Application Notes: | WB: 1.25 µg/mL<br>Optimal conditions should be determined by the investigator.   |
| Comment:           | TRIM17 Blocking Peptide, catalog no. 33R-7164, is also available for use as a blocking control in assays to test for specificity of this TRIM17 antibody |
| Restrictions:      | For Research Use only  |

## Handling

|                  |   |
|------------------|---|
| Format:          | Lyophilized   |
| Concentration:   | Lot specific  |
| Buffer:          | Lyophilized powder. Add 100 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. |
| Handling Advice: | Avoid repeated freeze/thaw cycles.  |
| Storage:         | 4 °C/-20 °C   |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C.                                     |

## Images



### Western Blotting

**Image 1.** Western Blot showing TRIM17 antibody used at a concentration of 1-2 µg/ml to detect its target protein.