

Datasheet for ABIN925936

anti-ZNF786 antibody (N-Term)

2 Images



| 0 | 100 | | | 100 | 100 | | | |
|----|-----|-------|---------|-----|----------|----|---|---|
| Go | †O | ν | \cap | dы | \cap t | na | | 0 |
| 00 | ιO | | \circ | uu | Uι | νu | ч | |

| _ | | | | | | |
|---|-----|----------|----|----|----|----|
| | 1// | Д | rv | 16 | ٦/ | ٨ |
| U | W | \vdash | ΙV | Ιt | ٦, | /V |

| Quantity: | 100 μg |
|----------------------|--|
| Target: | ZNF786 |
| Binding Specificity: | N-Term |
| Reactivity: | Human, Mouse, Rat, Dog, Cow, Chicken, Xenopus laevis |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ZNF786 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC) |
| Product Details | |
| Immunogen: | ZNF786 antibody was raised in rabbit using the N terminal of ZNF786 as the immunogen |
| Cross-Reactivity: | Mouse (Murine), Rat (Rattus), Cow (Bovine), Dog (Canine), Frog, Chicken |
| Target Details | |
| Target: | |
| raryet. | ZNF786 |
| Alternative Name: | ZNF786 ZNF786 (ZNF786 Products) |

Application Details

| Application Notes: | WB: 1.25 μg/mL | | |
|--------------------|--|--|--|
| | Optimal conditions should be determined by the investigator. | | |
| Comment: | ZNF786 Blocking Peptide, catalog no. 33R-5642, is also available for use as a blocking control in assays to test for specificity of this ZNF786 antibody | | |
| Restrictions: | For Research Use only | | |

Handling


| Format: | Lyophilized |
|------------------|--|
| Concentration: | Lot specific |
| Buffer: | Lyophilized powder. Add 100 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. |
| Handling Advice: | Avoid repeated freeze/thaw cycles. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C. |

Images



Western Blotting

Image 1. ZNF786 antibody (20R-1235) used at 0.2-1 ug/ml to detect target protein.



Immunohistochemistry

Image 2. ZNF786 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Alveolar cells (arrows) in Human Lung. Magnification is at 400X