

Datasheet for ABIN925968
anti-SNX29 antibody (N-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µg
Target:	SNX29 (RUNDC2A)
Binding Specificity:	N-Term
Reactivity:	Human, Cow, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNX29 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	RUNDC2 A antibody was raised in rabbit using the N terminal of RUNDC2 as the immunogen
Cross-Reactivity:	Rat (Rattus), Cow (Bovine)

Target Details

Target:	SNX29 (RUNDC2A)
Alternative Name:	RUNDC2A (RUNDC2A Products)
Background:	The function of this protein remains unknown. Synonyms: Polyclonal RUNDC2A antibody, Anti-RUNDC2A antibody, RUN domain containing 2A antibody, FLJ12363 antibody.


Application Details

Application Notes:	Optimal conditions should be determined by the investigator.
Comment:	RUNDC2A Blocking Peptide, catalog no. 33R-6436, is also available for use as a blocking control in assays to test for specificity of this RUNDC2A antibody
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 100 μ L of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images




Immunohistochemistry

Image 1. RUNDC2A antibody was used for immunohistochemistry at a concentration of 4-8 μ g/ml to stain Epithelial cells of intestinal villus (arrows) in Human Intestine. Magnification is at 400X



Immunohistochemistry

Image 2. RUNDC2A antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml. Magnification is at 400X



Western Blotting

Image 3. Western Blot showing RUNDC2A antibody used at a concentration of 5.0 ug/ml against HepG2 Cell Lysate