

Datasheet for ABIN926454  
**anti-MAGOH antibody (N-Term)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µL
Target:	MAGOH
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAGOH antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	MAGOH antibody was raised in rabbit using the N terminal of MAGOH as the immunogen
Purification:	Purified

## Target Details

Target:	MAGOH
Alternative Name:	MAGOH ( <a href="#">MAGOH Products</a> )
Background:	Drosophila that have mutations in their mago nashi (grandchildless) gene produce progeny with defects in germplasm assembly and germline development. MAGOH is the mammalian mago nashi homolog. In mammals, mRNA expression is not limited to the germ plasm, but is expressed ubiquitously in adult tissues and can be induced by serum stimulation of quiescent fibroblasts. Synonyms: Polyclonal MAGOH antibody, Anti-MAGOH antibody, mago-nashi

## Target Details

homolog, proliferation-associated, Drosophila antibody, MAGOH1 antibody, MAGOHA antibody.


## Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	MAGOH Blocking Peptide, catalog no. 33R-4537, is also available for use as a blocking control in assays to test for specificity of this MAGOH antibody
Restrictions:	For Research Use only

## Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

## Images



### Western Blotting

**Image 1.** MAGOH antibody (70R-10030) used at 0.2-1 µg/ml to detect target protein.