

Datasheet for ABIN926574  
**anti-HSH2D antibody (C-Term)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µL
Target:	HSH2D
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HSH2D antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	HSH2 D antibody was raised in rabbit using the C terminal of HSH2 as the immunogen
Purification:	Purified

## Target Details

Target:	HSH2D
Alternative Name:	HSH2D ( <a href="#">HSH2D Products</a> )
Background:	T-cell activation requires 2 signals: recognition of antigen by the T-cell receptor and a costimulatory signal provided primarily by CD28 in naive T cells. HSH2 is a target of both of these signaling pathways. Synonyms: Polyclonal HSH2D antibody, Anti-HSH2D antibody, hematopoietic SH2 domain containing antibody, ALX antibody, FLJ14886 antibody, HSH2 antibody.


## Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	HSH2D Blocking Peptide, catalog no. 33R-2096, is also available for use as a blocking control in assays to test for specificity of this HSH2D antibody
Restrictions:	For Research Use only

## Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

## Images



### Western Blotting

**Image 1.** Western Blot showing HSH2D antibody used at a concentration of 1-2 ug/ml to detect its target protein.