

Datasheet for ABIN926815
anti-TBL1XR1 antibody (C-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	TBL1XR1
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TBL1XR1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	TBL1 XR1 antibody was raised in rabbit using the C terminal of TBL1 R1 as the immunogen
Purification:	Purified

Target Details

Target:	TBL1XR1
Alternative Name:	TBL1XR1 (TBL1XR1 Products)
Background:	The protein encoded by this gene has sequence similarity with members of the WD40 repeat-containing protein family. The WD40 group is a large family of proteins, which appear to have a regulatory function. It is believed that the WD40 repeats mediate protein-protein interactions and members of the family are involved in signal transduction, RNA processing, gene regulation, vesicular trafficking, cytoskeletal assembly and may play a role in the control of

Target Details

cytotypic differentiation. Synonyms: Polyclonal TBL1XR1 antibody, Anti-TBL1XR1 antibody, transducin, beta-like 1 X-linked receptor 1 antibody, C21 antibody, DC42 antibody, FLJ12894 antibody, IRA1 antibody, TBLR1 antibody.

Pathways: [Regulation of Lipid Metabolism by PPARalpha](#)

Application Details

Application Notes: Optimal conditions should be determined by the investigator.

Comment: TBL1XR1 Blocking Peptide, catalog no. 33R-8490, is also available for use as a blocking control in assays to test for specificity of this TBL1XR1 antibody

Restrictions: For Research Use only

Handling

Format: Lyophilized

Concentration: Lot specific


Buffer: Lyophilized powder. Add 50 μ L of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: 4 °C/-20 °C

Storage Comment: Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. Western Blot showing TBL1XR1 antibody used at a concentration of 1 μ g/ml against Jurkat Cell Lysate