

Datasheet for ABIN926870  
**anti-TRADD antibody (Middle Region)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µL
Target:	TRADD
Binding Specificity:	Middle Region
Reactivity:	Human, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRADD antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	TRADD antibody was raised in rabbit using the middle region of TRADD as the immunogen
Cross-Reactivity:	Dog (Canine)
Purification:	Purified

## Target Details

Target:	TRADD
Alternative Name:	TRADD ( <a href="#">TRADD Products</a> )
Background:	<p>TRADD is the adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD.</p> <p>Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation of NF-kappa-B. The protein encoded by this gene is a death domain containing adaptor</p>

## Target Details

---

molecule that interacts with TNFRSF1A/TNFR1 and mediates programmed cell death signaling and NF-kappaB activation. Synonyms: Polyclonal TRADD antibody, Anti-TRADD antibody, TNFRSF1A-associated via death domain antibody, Hs.89862 antibody, MGC11078 antibody.

Pathways: [NF-kappaB Signaling](#), [Apoptosis](#), [Caspase Cascade in Apoptosis](#), [Positive Regulation of Endopeptidase Activity](#), [Hepatitis C](#)

## Application Details

---

Application Notes: WB: 0.2-1 µg/mL  
Optimal conditions should be determined by the investigator.

Comment: TRADD Blocking Peptide, catalog no. 33R-10088, is also available for use as a blocking control in assays to test for specificity of this TRADD antibody

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized


Concentration: Lot specific

Buffer: Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: 4 °C/-20 °C

Storage Comment: Store at 4 °C, following reconstitution, aliquot and store at -20 °C.



#### Western Blotting

**Image 1.** TRADD antibody (70R-10446) used at 0.2-1 ug/ml to detect target protein.