

## Datasheet for ABIN926889

# anti-DR3/LARD antibody (Middle Region)





Go to Product page

$\sim$			
( )\	<b>/</b> e	rVI	iew

O V CI V I C V V	
Quantity:	100 μL
Target:	DR3/LARD (TNFRSF25)
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DR3/LARD antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	TNFRSF25 antibody was raised in rabbit using the middle region of TNFRSF25 as the
	immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine)
Purification:	Purified
Target Details	
Target:	DR3/LARD (TNFRSF25)
Altanativa Nama	
Alternative Name:	TNFRSF25 (TNFRSF25 Products)
Background:	TNFRSF25 (TNFRSF25 Products)  TNFRSF25 is a member of the TNF-receptor superfamily. This receptor is expressed
	TNFRSF25 is a member of the TNF-receptor superfamily. This receptor is expressed

#### **Target Details**

regulate cell apoptosis. The signal transduction of this receptor is mediated by various death domain containing adaptor proteins. Knockout studies in mice suggested the role of this gene in the removal of self-reactive T cells in the thymus. Synonyms: Polyclonal TNFRSF25 antibody, Anti-TNFRSF25 antibody, tumor necrosis factor receptor superfamily, member 25 antibody, APO-3 antibody, DDR3 antibody, DR3 antibody, LARD antibody, TNFRSF12 antibody, TR3 antibody, TRAMP antibody, WSL-1 antibody, WSL-LR antibody.

Pathways:

**Apoptosis** 

### **Application Details**

Application Notes:	WB: 0.2-1 μg/mL		
	Optimal conditions should be determined by the investigator.		
Comment:	TNFRSF25 Blocking Peptide, catalog no. 33R-3528, is also available for use as a blocking control in assays to test for specificity of this TNFRSF25 antibody		
Restrictions:	For Research Use only		

#### Handling

Format:	Lyophilized	
Concentration:	Lot specific	
Buffer:	Lyophilized powder. Add 50 $\mu$ L of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.	
Handling Advice:	Avoid repeated freeze/thaw cycles.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.	

90 kDa\_\_ 65 kDa\_\_ 40 kDa\_\_ 31 kDa\_\_ 22 kDa\_\_

### **Western Blotting**

**Image 1.** TNFRSF25 antibody (70R-10465) used at 0.2-1 ug/ml to detect target protein.