

Datasheet for ABIN927274
anti-TOX2 antibody (N-Term)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	TOX2
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Dog, Chicken, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TOX2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	C20 ORF100 antibody was raised in rabbit using the N terminal of C20 RF100 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Dog (Canine), Chicken, Frog
Purification:	Purified

Target Details

Target:	TOX2
Alternative Name:	C20ORF100 (TOX2 Products)
Background:	This gene, which is located on chromosome 20, is predicted to encode for a signal protein with unknown function. Synonyms: Polyclonal C20ORF100 antibody, Anti-C20ORF100 antibody.


Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	C20ORF100 Blocking Peptide, catalog no. 33R-9294, is also available for use as a blocking control in assays to test for specificity of this C20ORF100 antibody
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Immunohistochemistry

Image 1. C20ORF100 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Epithelial cells of intestinal villus (arrows) in Human Intestine. Magnification is at 400X



Western Blotting

Image 2. C20ORF100 antibody used at 0.1-2 ug/ml on a Jurkat cell lysate.