

Datasheet for ABIN927334
anti-FGD1 antibody (C-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	FGD1
Binding Specificity:	C-Term
Reactivity:	Human, Cow, Dog, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FGD1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	FGD1 antibody was raised in rabbit using the C terminal of FGD1 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine), Dog (Canine)
Purification:	Purified

Target Details

Target:	FGD1
Alternative Name:	FGD1 (FGD1 Products)
Background:	FGD1 contains Dbl (DH) and pleckstrin (PH) homology domains. It can bind specifically to the Rho family GTPase Cdc42Hs and stimulate the GDP-GTP exchange of the isoprenylated form of Cdc42Hs. It also stimulates the mitogen activated protein kinase cascade leading to c-Jun kinase SAPK/JNK1 activation. FGD1 has an essential role in embryonic development, and FGD1

Target Details

gene mutations result in the human developmental disorder, Aarskog-Scott syndrome.

Synonyms: Polyclonal FGD1 antibody, Anti-FGD1 antibody, FYVE, RhoGEF and PH domain containing 1, faciogenital dysplasia antibody.

Pathways: [Neurotrophin Signaling Pathway](#)

Application Details

Application Notes: WB: 0.2-1 µg/mL
Optimal conditions should be determined by the investigator.

Comment: FGD1 Blocking Peptide, catalog no. 33R-9977, is also available for use as a blocking control in assays to test for specificity of this FGD1 antibody

Restrictions: For Research Use only

Handling

Format: Lyophilized

Concentration: Lot specific

Buffer: Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: 4 °C/-20 °C

Storage Comment: Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. FGD1 antibody used at 0.2-1 ug/ml to detect target protein.