

Datasheet for ABIN927338 **anti-ZNF385A antibody (C-Term)**



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	ZNF385A
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Cow, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF385A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	ZNF385 antibody was raised in rabbit using the C terminal of ZNF385 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Zebrafish (Brachydanio rerio), Cow (Bovine)
Purification:	Purified

Target Details

Target:	ZNF385A
Alternative Name:	ZNF385 (ZNF385A Products)
Background:	Results suggest that RZF is a shuttling regulatory protein expressed in photoreceptors of the human retina that may be involved in mRNA or protein regulation of photoreceptor-specific genes and therefore have role in retinal disease mechanisms. Synonyms: Polyclonal ZNF385 antibody, Anti-ZNF385 antibody, zinc finger protein 385 antibody.

Target Details

Pathways: [Positive Regulation of Response to DNA Damage Stimulus](#), [Negative Regulation of intrinsic apoptotic Signaling](#), [Positive Regulation of fat Cell Differentiation](#)

Application Details

Application Notes: WB: 0.2-1 µg/mL
Optimal conditions should be determined by the investigator.

Comment: ZNF385 Blocking Peptide, catalog no. 33R-5098, is also available for use as a blocking control in assays to test for specificity of this ZNF385 antibody

Restrictions: For Research Use only

Handling

Format: Lyophilized

Concentration: Lot specific


Buffer: Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: 4 °C/-20 °C


Storage Comment: Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. ZNF385 antibody used at 0.2-1 µg/ml to detect target protein.



Immunohistochemistry

Image 2. ZNF385 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Epidermal cells (arrows) in Human Skin. Magnification is at 400X