

Datasheet for ABIN927339

anti-ZBTB32 antibody (N-Term)

1 Image



Go to Product page

| \sim | | | | |
|--------|-----|-----|-----|---|
| () | ve. | r\/ | 101 | Λ |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | ZBTB32 | |
| Binding Specificity: | N-Term | |
| Reactivity: | Human, Mouse, Cow, Dog | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This ZBTB32 antibody is un-conjugated | |
| Application: | Western Blotting (WB) | |
| Product Details | | |
| Immunogen: | ZBTB32 antibody was raised in rabbit using the N terminal of ZBTB32 as the immunogen | |
| Cross-Reactivity: | Dog (Canine), Mouse (Murine), Cow (Bovine) | |
| Purification: | Purified | |
| Target Details | | |
| Target: | ZBTB32 | |
| Alternative Name: | ZBTB32 (ZBTB32 Products) | |
| Background: | ZBTB32 may play an essential role during the proliferative stages of primitive hematopoietic progenitors, possibly acting in concert with (a subset of) the Fanconi anemia proteins. This gene can also interact with GATA-2 and can modify GATA-2 transactivation capacity. Synonyms: Polyclonal ZBTB32 antibody, Anti-ZBTB32 antibody, zinc finger and BTB domain | |

| containing | 32 | antibo | dy |
|------------|----|--------|----|
|------------|----|--------|----|


Application Details

| Application Notes: | WB: 0.2-1 μg/mL |
|--------------------|--|
| | Optimal conditions should be determined by the investigator. |
| Comment: | ZBTB32 Blocking Peptide, catalog no. 33R-7162, is also available for use as a blocking control in assays to test for specificity of this ZBTB32 antibody |
| Restrictions: | For Research Use only |

Handling

| Format: | Lyophilized | |
|------------------|---|--|
| Concentration: | Lot specific | |
| Buffer: | Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. | |
| Handling Advice: | Avoid repeated freeze/thaw cycles. | |
| Storage: | 4 °C/-20 °C | |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C. | |

Images



Western Blotting

Image 1. ZBTB32 antibody used at 0.2-1 ug/ml to detect target protein.