

Datasheet for ABIN927377
anti-DACH2 antibody (C-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	DACH2
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Dog, Zebrafish (Danio rerio), Chicken, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DACH2 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	DACH2 antibody was raised in rabbit using the C terminal of DACH2 as the immunogen
Cross-Reactivity:	Mouse (Murine), Zebrafish (Brachydanio rerio), Frog, Chicken, Dog (Canine)
Purification:	Purified

Target Details

Target:	DACH2
Alternative Name:	DACH2 (DACH2 Products)
Background:	Most X autosome translocations associated with premature ovarian failure do not interrupt X-linked genes. Only one of the six breakpoints disrupts the DACH2 gene. Synonyms: Polyclonal DACH2 antibody, Anti-DACH2 antibody, dachshund homolog 2, Drosophila antibody.

Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	DACH2 Blocking Peptide, catalog no. 33R-9147, is also available for use as a blocking control in assays to test for specificity of this DACH2 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. DACH2 antibody used at 0.2-1 ug/ml to detect target protein.