

Datasheet for ABIN927393

anti-ZNF385A antibody (N-Term)

2 Images



Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Quantity:	100 μL
Target:	ZNF385A
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF385A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Troduct Details	
Immunogen:	ZNF385 antibody was raised in rabbit using the N terminal of ZNF385 as the immunogen
	ZNF385 antibody was raised in rabbit using the N terminal of ZNF385 as the immunogen Mouse (Murine), Rat (Rattus), Cow (Bovine)
lmmunogen:	
Immunogen: Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine)
Immunogen: Cross-Reactivity: Purification:	Mouse (Murine), Rat (Rattus), Cow (Bovine)
Immunogen: Cross-Reactivity: Purification: Target Details	Mouse (Murine), Rat (Rattus), Cow (Bovine) Purified
Immunogen: Cross-Reactivity: Purification: Target Details Target:	Mouse (Murine), Rat (Rattus), Cow (Bovine) Purified ZNF385A

apoptotic Signaling, Positive Regulation of fat Cell Differentiation


Application Details

Application Notes:	WB: 0.2-1 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	ZNF385 Blocking Peptide, catalog no. 33R-2624, is also available for use as a blocking control in assays to test for specificity of this ZNF385 antibody
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Immunohistochemistry

Image 1. ZNF385 antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml to stain Alveolar ceils (arrows) in Human Lung. Magnification is at 400X



Western Blotting

Image 2. ZNF385 antibody used at 0.2-1 ug/ml to detect target protein.