

# Datasheet for ABIN927423 anti-CHRAC1 antibody (N-Term)

## 1 Image



Go	to	Pro	du	ıct	na	76
GU	ιΟ	FIL	uu	IUL	pa	45

0				

Overview	
Quantity:	100 μL
Target:	CHRAC1
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CHRAC1 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	CHRAC1 antibody was raised in rabbit using the N terminal of CHRAC1 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine)
Purification:	Purified
Target Details	
Target:	CHRAC1
Alternative Name:	CHRAC1 (CHRAC1 Products)
Background:	CHRAC1 is a histone-fold protein that interacts with other histone-fold proteins to bind DNA in a sequence-independent manner. These histone-fold protein dimers combine within larger enzymatic complexes for DNA transcription, replication, and packaging. CHRAC1 is a histone-

fold protein that interacts with other histone-fold proteins to bind DNA in a sequence-

independent manner. These histone-fold protein dimers combine within larger enzymatic complexes for DNA transcription, replication, and packaging. Synonyms: Polyclonal CHRAC1 antibody, Anti-CHRAC1 antibody, chromatin accessibility complex 1 antibody, CHARC1 antibody, CHARC15 antibody, YCL1 antibody.


### **Application Details**

Application Notes:	WB: 0.2-1 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	CHRAC1 Blocking Peptide, catalog no. 33R-8534, is also available for use as a blocking control in assays to test for specificity of this CHRAC1 antibody
Restrictions:	For Research Use only

### Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 $\mu L$ of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

#### **Images**



#### **Western Blotting**

**Image 1.** CHRAC1 antibody used at 1.25 ug/ml to detect target protein.