

Datasheet for ABIN927426  
**anti-FA2H antibody (C-Term)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µL
Target:	FA2H
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FA2H antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	Fa2 h antibody was raised in rabbit using the C terminal of Fa2 as the immunogen
Cross-Reactivity:	Mouse (Murine)
Purification:	Purified

## Target Details

Target:	FA2H
Alternative Name:	Fa2h ( <a href="#">FA2H Products</a> )
Background:	Fa2h is required for alpha-hydroxylation of free fatty acids and the formation of alpha-hydroxylated sphingolipids. Synonyms: Polyclonal Fa2h antibody, Anti-Fa2h antibody, fatty acid 2-hydroxylase antibody, FAAH antibody, Faxdc1 antibody, G630055L08Rik antibody.


## Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	Fa2h Blocking Peptide, catalog no. 33R-3743, is also available for use as a blocking control in assays to test for specificity of this Fa2h antibody
Restrictions:	For Research Use only

## Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

## Images



### Western Blotting

**Image 1.** Fa2h antibody used at 2.5 µg/ml to detect target protein.