

Datasheet for ABIN927442 anti-STAT5B antibody (N-Term)





\sim				
()	ve.	r\/	101	Λ

100 μL	
STAT5B	
N-Term	
Human, Mouse, Rat, Cow, Dog, Pig, Chicken	
Rabbit	
Polyclonal	
This STAT5B antibody is un-conjugated	
Western Blotting (WB)	
STAT5 B antibody was raised in rabbit using the N terminal of STAT5 as the immunogen	
Mouse (Murine), Rat (Rattus), Pig (Porcine), Dog (Canine), Cow (Bovine), Chicken	
Mouse (Murine), Rat (Rattus), Pig (Porcine), Dog (Canine), Cow (Bovine), Chicken	
Mouse (Murine), Rat (Rattus), Pig (Porcine), Dog (Canine), Cow (Bovine), Chicken Purified	
Purified	

Target Details

various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. Synonyms: Polyclonal STAT5B antibody, Anti-STAT5B antibody, signal transducer and activator of transcription 5B antibody, STAT5 antibody.

Pathways:


JAK-STAT Signaling, RTK Signaling, Response to Growth Hormone Stimulus, C21-Steroid Hormone Metabolic Process, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, CXCR4-mediated Signaling Events, Activated T Cell Proliferation

Application Details

Application Notes:	Optimal conditions should be determined by the investigator.	
Comment:	STAT5B Blocking Peptide, catalog no. 33R-5362, is also available for use as a blocking control in assays to test for specificity of this STAT5B antibody	
Restrictions:	For Research Use only	


Handling

Format:	Lyophilized	
Concentration:	Lot specific	
Buffer:	Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.	
Handling Advice:	Avoid repeated freeze/thaw cycles.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.	



Immunohistochemistry

Image 1. STAT5B antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml. Magnification is at 400X



Immunohistochemistry

Image 2. STAT5B antibody was used for immunohistochemistry at a concentration of 4-8 ug/ml. Magnification is at 400X

90 kDa__ 65 kDa__ 40 kDa__ 31 kDa__ 22 kDa__

Western Blotting

Image 3. Western Blot showing STAT5B antibody used at a concentration of 1.0 ug/ml against Transfected 293T Cell Lysate