

Datasheet for ABIN927491 **anti-PHD1 antibody (N-Term)**



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	PHD1 (EGLN2)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PHD1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	EGLN2 antibody was raised in rabbit using the N terminal of EGLN2 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Dog (Canine)
Purification:	Purified

Target Details

Target:	PHD1 (EGLN2)
Alternative Name:	EGLN2 (EGLN2 Products)
Background:	The hypoxia inducible factor (HIF) is a transcriptional complex which is involved in oxygen homeostasis. At normal oxygen levels, the alpha subunit of HIF is targeted for degradation by prolyl hydroxylation. EGLN2 encodes an enzyme responsible for this posttranslational modification. Alternative splicing of EGLN2 results in three transcript variants encoding

Target Details

different isoforms. Synonyms: Polyclonal EGLN2 antibody, Anti-EGLN2 antibody, egl nine homolog 2, C. elegans antibody.

Pathways: [Intracellular Steroid Hormone Receptor Signaling Pathway](#), [Cell RedoxHomeostasis](#)

Application Details

Application Notes: WB: 0.2-1 µg/mL
Optimal conditions should be determined by the investigator.

Comment: EGLN2 Blocking Peptide, catalog no. 33R-8295, is also available for use as a blocking control in assays to test for specificity of this EGLN2 antibody

Restrictions: For Research Use only

Handling

Format: Lyophilized

Concentration: Lot specific


Buffer: Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: 4 °C/-20 °C

Storage Comment: Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. EGLN2 antibody used at 1.25 µg/ml to detect target protein.