

Datasheet for ABIN927496

anti-TRIM17 / RNF16 antibody (Middle Region)[Go to Product page](#)**2** Images

Overview

Quantity:	100 µL
Target:	TRIM17 / RNF16 (TRIM17)
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM17 / RNF16 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	TRIM17 antibody was raised in rabbit using the middle region of TRIM17 as the immunogen
Cross-Reactivity:	Rat (Rattus), Mouse (Murine), Cow (Bovine)
Purification:	Purified

Target Details

Target:	TRIM17 / RNF16 (TRIM17)
Alternative Name:	TRIM17 (TRIM17 Products)
Background:	TRIM17 encodes a protein that is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic bodies. The protein is expressed at high levels in the testis, but its function is unknown. Synonyms: Polyclonal TRIM17 antibody, Anti-

Target Details

TRIM17 antibody, tripartite motif-containing 17 antibody.


Application Details

Application Notes:	WB: 0.12 µg/mL Optimal conditions should be determined by the investigator.
Comment:	TRIM17 Blocking Peptide, catalog no. 33R-6134, is also available for use as a blocking control in assays to test for specificity of this TRIM17 antibody
Restrictions:	For Research Use only

Handling


Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Immunohistochemistry

Image 1. TRIM17 antibody was used for immunohistochemistry at a concentration of 4-8 µg/ml to stain Liver cells (arrows) in Human Liver. Magnification is at 400X



Western Blotting

Image 2. TRIM17 antibody used at 2.5 ug/ml to detect target protein.