

Datasheet for ABIN927516 anti-DIDO1 antibody (N-Term)

1 Image



Go to Product page


_						
	1//	Д	rv	16	٦/	٨
U	W	\vdash	ΙV	Ιt	٦,	/V

Overview	
Quantity:	100 μL
Target:	DID01
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DIDO1 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	DIDO1 antibody was raised in rabbit using the N terminal of DIDO1 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine), Chicken
Purification:	Purified
Target Details	
Target:	DID01
Alternative Name:	DID01 (DID01 Products)
Background:	In mice, the death inducer-obliterator-1 gene is upregulated by apoptotic signals and encodes a
	cytoplasmic protein that translocates to the nucleus upon apoptotic signal activation. When
	overexpressed, the mouse protein induced apoptosis in cell lines growing in vitro. DIDO1 gene
	is similar to the mouse gene and therefore is thought to be involved in apoptosis. Synonyms:

Target Details

l arget Details	
	Polyclonal DID01 antibody, Anti-DID01 antibody, death inducer-obliterator 1 antibody.
Pathways:	Hormone Activity
Application Details	
Application Notes:	WB: 0.2-1 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	DIDO1 Blocking Peptide, catalog no. 33R-2714, is also available for use as a blocking control in
	assays to test for specificity of this DIDO1 antibody
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in
	PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. DIDO1 antibody used at 2.5 ug/ml to detect target protein.