

## Datasheet for ABIN927601

# anti-Zinc Finger Protein 3 Homolog (ZFP3) (N-Term) antibody





Go to Product page

| $\sim$ |     |       |    |
|--------|-----|-------|----|
|        | ve  | r\/   | ٨  |
| $\cup$ | V C | 1 V I | ٧V |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | Zinc Finger Protein 3 Homolog (ZFP3)  |  |
| Binding Specificity: | N-Term  |  |
| Reactivity:          | Human   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | Un-conjugated   |  |
| Application:         | Western Blotting (WB)   |  |
| Product Details      |   |  |
| Immunogen:           | ZFP3 antibody was raised in rabbit using the N terminal of ZFP3 as the immunogen  |  |
| Purification:        | Purified  |  |
| Target Details       |   |  |
| Target:              | Zinc Finger Protein 3 Homolog (ZFP3)  |  |
| Alternative Name:    | ZFP3 (ZFP3 Products)  |  |
| Background:          | ZFP3 belongs to the krueppel C2H2-type zinc-finger protein family. It contains 13 C2H2-type zinc fingers. ZFP3 may be involved in transcriptional regulation. Synonyms: Polyclonal ZFP3 |  |

## **Application Details**

| Application Notes: | WB: 0.2-1 μg/mL  |  |
|--------------------|--|--|
|                    | Optimal conditions should be determined by the investigator.   |  |
| Comment:           | ZFP3 Blocking Peptide, catalog no. 33R-9312, is also available for use as a blocking control in assays to test for specificity of this ZFP3 antibody |  |
| Restrictions:      | For Research Use only  |  |

## Handling

| Format:          | Lyophilized   |  |
|------------------|---|--|
| Concentration:   | Lot specific  |  |
| Buffer:          | Lyophilized powder. Add 50 $\mu$ L of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. |  |
| Handling Advice: | Avoid repeated freeze/thaw cycles.  |  |
| Storage:         | 4 °C/-20 °C   |  |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C.   |  |

#### **Images**



#### **Western Blotting**

**Image 1.** ZFP3 antibody used at 0.2-1 ug/ml to detect target protein.