

Datasheet for ABIN927745
anti-WDHD1 antibody (C-Term)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	WDHD1
Binding Specificity:	C-Term
Reactivity:	Human, Cow, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This WDHD1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	WDHD1 antibody was raised in rabbit using the C terminal of WDHD1 as the immunogen
Cross-Reactivity:	Cow (Bovine), Dog (Canine)
Purification:	Purified

Target Details

Target:	WDHD1
Alternative Name:	WDHD1 (WDHD1 Products)
Background:	WDHD1 contains multiple N-terminal WD40 domains and a C-terminal high mobility group (HMG) box. WD40 domains are found in a variety of eukaryotic proteins and may function as adaptor/regulatory modules in signal transduction, pre-mRNA processing and cytoskeleton assembly. HMG boxes are found in many eukaryotic proteins involved in chromatin assembly,

Target Details

transcription and replication. WDHD1 may also be involved in protein-protein interactions.
Synonyms: Polyclonal WDHD1 antibody, Anti-WDHD1 antibody, WD repeat and HMG-box DNA binding protein 1 antibody, AND-1 antibody.

Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	WDHD1 Blocking Peptide, catalog no. 33R-4628, is also available for use as a blocking control in assays to test for specificity of this WDHD1 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.