

Datasheet for ABIN927794 anti-ZNF415 antibody (N-Term)

and Zivi 410 and body (iv 1





()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Quantity:	100 μL	
Target:	ZNF415	
Binding Specificity:	N-Term	
Reactivity:	Human, Rat, Mouse, Cow, Dog	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZNF415 antibody is un-conjugated	
Application:	Western Blotting (WB)	
Product Details		
Immunogen:	ZNF415 antibody was raised in rabbit using the N terminal of ZNF415 as the immunogen	
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Dog (Canine), Cow (Bovine)	
Purification:	Purified	
Target Details		
Target:	ZNF415	
Alternative Name:	ZNF415 (ZNF415 Products)	
Background:	ZNF415 is involved in transcriptional regulation. The transcriptional activity differed among the various isoforms. All isoforms except isoform 3 seem to suppress the transcriptional activities of AP-1 and p53. Synonyms: Polyclonal ZNF415 antibody, Anti-ZNF415 antibody, zinc finger	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn | International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com | Page 1/2 | Product datasheet for ABIN927794 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

protein 415 antibody, FLJ11191 antibody.


Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	ZNF415 Blocking Peptide, catalog no. 33R-5655, is also available for use as a blocking control in assays to test for specificity of this ZNF415 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized	
Concentration:	Lot specific	
Buffer:	Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.	
Handling Advice:	Avoid repeated freeze/thaw cycles.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.	

Images



Western Blotting

Image 1. ZNF415 antibody used at 5 ug/ml to detect target protein.