

Datasheet for ABIN927822
anti-ZNF335 antibody (Middle Region)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	ZNF335
Binding Specificity:	Middle Region
Reactivity:	Human, Rat, Cow, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF335 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	ZNF335 antibody was raised in rabbit using the middle region of ZNF335 as the immunogen
Cross-Reactivity:	Rat (Rattus), Cow (Bovine), Dog (Canine)
Purification:	Purified

Target Details

Target:	ZNF335
Alternative Name:	ZNF335 (ZNF335 Products)
Background:	ZNF335 enhances transcriptional activation by ligand-bound nuclear hormone receptors. However, it does this not by direct interaction with the receptor, but by direct interaction with the nuclear hormone receptor transcriptional coactivator NRC. ZNF335 may function by altering local chromatin structure. The protein encoded by this gene enhances transcriptional

Target Details


activation by ligand-bound nuclear hormone receptors. However, it does this not by direct interaction with the receptor, but by direct interaction with the nuclear hormone receptor transcriptional coactivator NRC. The encoded protein may function by altering local chromatin structure. Synonyms: Polyclonal ZNF335 antibody, Anti-ZNF335 antibody, zinc finger protein 335 antibody, NIF1 antibody.

Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	ZNF335 Blocking Peptide, catalog no. 33R-2248, is also available for use as a blocking control in assays to test for specificity of this ZNF335 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.



Western Blotting

Image 1. ZNF335 antibody used at 0.2-1 ug/ml to detect target protein.