ANTIBODIES ONLINE

Datasheet for ABIN927930 anti-U2AF2 antibody (C-Term)

Image



o to i roddot page

Overview

1

Quantity:	100 µL
Target:	U2AF2 (U2AF59)
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This U2AF2 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	U2 AF2 antibody was raised in rabbit using the C terminal of U2 F2 as the immunogen
Purification:	Purified
Target Details	
Target:	U2AF2 (U2AF59)
Alternative Name:	U2AF2 (U2AF59 Products)
Background:	U2 auxiliary factor (U2AF), comprised of a large and a small subunit, is a non-snRNP protein
	required for the binding of U2 snRNP to the pre-mRNA branch site. U2AF2 is the U2AF large
	subunit which contains a sequence-specific RNA-binding region with 3 RNA recognition motifs
	and an Arg/Ser-rich domain necessary for splicing. The large subunit binds to the
	polypyrimidine tract of introns early during spliceosome assembly. Synonyms: Polyclonal
Order et united	antibadias anlina com Luunu antikaarnar anlina da Luunu antiaarna anligna fr Luunu antibadias anlina an

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN927930 | 07/26/2024 | Copyright antibodies-online. All rights reserved.


Target Details

U2AF2 antibody, Anti-U2AF2 antibody, U2 small nuclear RNA auxiliary factor 2 antibody, U2AF65 antibody.

Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. U2AF2 antibody used at 0.2-1 ug/ml to detect target protein.