

Datasheet for ABIN927933

anti-AUH antibody (N-Term)





Go to Product page

_						
	1//	Д	rv	16	٦/	٨
U	W	\vdash	ΙV	Ιt	٦,	/V

Quantity:	100 μL
Target:	AUH
Binding Specificity:	N-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AUH antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	Auh antibody was raised in rabbit using the N terminal of Auh as the immunogen
Cross-Reactivity:	Mouse (Murine)
Purification:	Purified
Target Details	
Target:	AUH
Alternative Name:	Auh (AUH Products)
Background:	Auh Catalyzes the conversion of 3-methylglutaconyl-CoA to 3-hydroxy-3-methylglutaryl-CoA. Auh has very low enoyl-CoA hydratase activity. Was originally identified as RNA-binding protein

Auh antibody, AU RNA binding protein/enoyl-coenzyme A hydratase antibody, C77140 antibody,

W91705	antibody
--------	----------


Application Details

Application Notes:	WB: 0.2-1 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	Auh Blocking Peptide, catalog no. 33R-7340, is also available for use as a blocking control in assays to test for specificity of this Auh antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 μ L of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. Auh antibody used at 0.2-1 ug/ml to detect target protein.