

Datasheet for ABIN927944

anti-UTP18 antibody (N-Term)

1 Image




Overview

| Overview | |
|----------------------|---|
| Quantity: | 100 μL |
| Target: | UTP18 |
| Binding Specificity: | N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This UTP18 antibody is un-conjugated |
| Application: | Western Blotting (WB) |
| Product Details | |
| Immunogen: | UTP18 antibody was raised in rabbit using the N terminal of UTP18 as the immunogen |
| Purification: | Purified |
| Target Details | |
| Target: | UTP18 |
| Alternative Name: | UTP18 (UTP18 Products) |
| Background: | UTP18 is involved in nucleolar processing of pre-18S ribosomal RNA. Synonyms: Polyclonal UTP18 antibody, Anti-UTP18 antibody, UTP18, small subunit, SSU processome component, homolog, yeast antibody, CGI-48 antibody, WDR50 antibody. |

Application Details

| Application Notes: | WB: 0.2-1 μ g/mL Optimal conditions should be determined by the investigator. |
|--------------------|---|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Lyophilized |
| Concentration: | Lot specific |
| Buffer: | Lyophilized powder. Add 50 μ L of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer. |
| Handling Advice: | Avoid repeated freeze/thaw cycles. |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Store at 4 °C, following reconstitution, aliquot and store at -20 °C. |

Images



Western Blotting

Image 1. UTP18 antibody used at 0.2-1 ug/ml to detect target protein.