

Datasheet for ABIN928004  
**anti-NR2F1 antibody (N-Term)**



[Go to Product page](#)

1 Image

## Overview

Quantity:	100 µL
Target:	NR2F1
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Zebrafish (Danio rerio), Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR2F1 antibody is un-conjugated
Application:	Western Blotting (WB)

## Product Details

Immunogen:	NR2 F1 antibody was raised in rabbit using the N terminal of NR2 1 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine), Frog, Zebrafish (Brachydanio rerio)
Purification:	Purified

## Target Details

Target:	NR2F1
Alternative Name:	NR2F1 ( <a href="#">NR2F1 Products</a> )
Background:	Coup (chicken ovalbumin upstream promoter) transcription factor binds to the ovalbumin promoter and, in conjunction with another protein (S300-II) stimulates initiation of transcription. NR2F1 binds to both direct repeats and palindromes of the 5'-AGGTCA-3' motif. Synonyms: Polyclonal NR2F1 antibody, Anti-NR2F1 antibody, nuclear receptor subfamily 2, group F,

## Target Details

member 1 antibody, COUP-TFI antibody, EAR-3 antibody, EAR3 antibody, ERBAL3 antibody, NR2F2 antibody, SVP44 antibody, TCFCOUP1 antibody, TFCOUP1 antibody.

Pathways: [Nuclear Receptor Transcription Pathway](#), [Steroid Hormone Mediated Signaling Pathway](#)

## Application Details

Application Notes: WB: 0.2-1 µg/mL  
Optimal conditions should be determined by the investigator.

Comment: NR2F1 Blocking Peptide, catalog no. 33R-1391, is also available for use as a blocking control in assays to test for specificity of this NR2F1 antibody

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Concentration: Lot specific

Buffer: Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: 4 °C/-20 °C

Storage Comment: Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

## Images



### Western Blotting

**Image 1.** NR2F1 antibody used at 0.2-1 ug/ml to detect target protein.