

Datasheet for ABIN928150
anti-ISX antibody (N-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	ISX
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ISX antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	ISX antibody was raised in rabbit using the N terminal of ISX as the immunogen
Purification:	Purified

Target Details

Target:	ISX
Alternative Name:	ISX (ISX Products)
Background:	Homeobox genes encode DNA-binding proteins, many of which are thought to be involved in early embryonic development. Homeobox genes encode a DNA-binding domain of 60 to 63 amino acids referred to as the homeodomain. This gene is a member of the RAXLX homeobox gene family. Homeobox genes encode DNA-binding proteins, many of which are thought to be involved in early embryonic development. Homeobox genes encode a DNA-binding domain of

Target Details

60 to 63 amino acids referred to as the homeodomain. This gene is a member of the RAXLX homeobox gene family. Synonyms: Polyclonal ISX antibody, Anti-ISX antibody, intestine-specific homeobox antibody, DKFZp781N2395 antibody, MGC138417 antibody, RAXLX antibody.


Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	ISX Blocking Peptide, catalog no. 33R-4053, is also available for use as a blocking control in assays to test for specificity of this ISX antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. ISX antibody used at 0.2-1 ug/ml to detect target protein.