

Datasheet for ABIN928173 **anti-TAF13 antibody (N-Term)**



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	TAF13
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Cow, Dog, Chicken, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAF13 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	TAF13 antibody was raised in rabbit using the N terminal of TAF13 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Cow (Bovine), Dog (Canine), Chicken, Frog
Purification:	Purified

Target Details

Target:	TAF13
Alternative Name:	TAF13 (TAF13 Products)
Background:	TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. TAF13 is a small subunit

Target Details

associated with a subset of TFIID complexes. TAF13 interacts with TBP and with two other small subunits of TFIID, TAF10 and TAF11. Synonyms: Polyclonal TAF13 antibody, Anti-TAF13 antibody, TAF13 RNA polymerase II, TATA box binding protein, TBP-associated factor, 18kDa antibody, MGC22425 antibody, TAF2K antibody, TAFII18 antibody.


Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	TAF13 Blocking Peptide, catalog no. 33R-1087, is also available for use as a blocking control in assays to test for specificity of this TAF13 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. TAF13 antibody used at 0.2-1 ug/ml to detect target protein.