

Datasheet for ABIN928429
anti-Cyclin L2 antibody (N-Term)



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	Cyclin L2 (CCNL2)
Binding Specificity:	N-Term
Reactivity:	Human, Rat, Mouse, Dog, Chicken, Xenopus laevis, Drosophila melanogaster
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cyclin L2 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	CCNL2 antibody was raised in rabbit using the N terminal of CCNL2 as the immunogen
Cross-Reactivity:	Mouse (Murine), Rat (Rattus), Chicken, Frog, Dog (Canine), Fruit Fly (Drosophila melanogaster)
Purification:	Purified

Target Details

Target:	Cyclin L2 (CCNL2)
Alternative Name:	CCNL2 (CCNL2 Products)
Background:	CCNL2 belongs to the cyclin family, Cyclin L subfamily. CCNL2 is a transcriptional regulator which participates in regulating the pre-mRNA splicing process. CCNL2 also modulates the expression of critical apoptotic factor, leading to cell apoptosis. Synonyms: Polyclonal CCNL2 antibody, Anti-CCNL2 antibody, cyclin L2 antibody, ANIA-6B antibody, DKFZp761A1210

Target Details

antibody, DKFZp7620195 antibody, HCLA-ISO antibody, HLA-ISO antibody, PCEE antibody, SB138 antibody.

Application Details

Application Notes:	WB: 0.2-1 µg/mL Optimal conditions should be determined by the investigator.
Comment:	CCNL2 Blocking Peptide, catalog no. 33R-9094, is also available for use as a blocking control in assays to test for specificity of this CCNL2 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 µL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. CCNL2 antibody used at 0.2-1 ug/ml to detect target protein.