

Datasheet for ABIN928455 anti-ZNF781 antibody (N-Term)

1 Image



Overview

Quantity:	100 μL
Target:	ZNF781
Binding Specificity:	N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB)

Product Details

Immunogen:	ZNF781 antibody was raised in rabbit using the N terminal of ZNF781 as the immunogen
Purification:	Purified

Target Details

Target:	ZNF /81
Alternative Name:	ZNF781 (ZNF781 Products)
Background:	ZNF781 belongs to the krueppel C2H2-type zinc-finger protein family. It contains 4 C2H2-type

ZNF781 belongs to the krueppel C2H2-type zinc-finger protein family. It contains 4 C2H2-type zinc fingers. ZNF781 may be involved in transcriptional regulation. The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). Synonyms: Polyclonal ZNF781 antibody, Anti-ZNF781 antibody, zinc finger protein 781 antibody, FLJ37549

antibody, MGC131783 antibody	antibody.	MGC1	31783	antibody	٧.
------------------------------	-----------	------	-------	----------	----


Application Details

Application Notes:	WB: 0.2-1 μg/mL
	Optimal conditions should be determined by the investigator.
Comment:	ZNF781 Blocking Peptide, catalog no. 33R-7715, is also available for use as a blocking control in assays to test for specificity of this ZNF781 antibody
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Concentration:	Lot specific
Buffer:	Lyophilized powder. Add 50 μL of distilled water. Final antibody concentration is 1 mg/mL in PBS buffer.
Handling Advice:	Avoid repeated freeze/thaw cycles.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 4 °C, following reconstitution, aliquot and store at -20 °C.

Images



Western Blotting

Image 1. ZNF781 antibody used at 1.25 ug/ml to detect target protein.